

ANTARCTIC GEOLOGICAL 1:250,000 MAP SERIES  
MOUNT JOYCE QUADRANGLE  
(VICTORIA LAND)  
1999

G. Capponi<sup>1</sup>, L. Crispini<sup>2</sup>, M. Meccheri<sup>2</sup>, G. Musumeci<sup>2</sup> & P.C. Pertusati<sup>3</sup>



Authors:  
C. Baroni<sup>4</sup>, C. Delisle<sup>4</sup>, G. Orsi<sup>5</sup>

- Scree, glacial debris and moraine (n). Outwash.
  - Ricker Hills tillite (Rt): massive matrix-supported diamictite (Sirius Formation?).
- POST-ROSS MAGMATISM AND SEDIMENTATION**
- KIRKPATRICK BASALT (Kb)**  
Subspherical, a few metres up to several ten metres thick, randomly separated by thin silty sedimentary volcanoclastic interlayers and pillow lavas. Blue lines (omitted) show the original, vertical, columnar jointing of which many striae, with traua and rock relics mostly in the lower portion of the volcanic sequence.
  - FERRAR DOLERITE (Fd)**  
Tholeiitic dolerite sills and minor dykes, usually intruded in the sedimentary sequence of the Beacon Supergroup, immediately above the pre-Beacon peneplain. Black lines (omitted) in the cross-sectional maps denote and separate (nearly some ten metres thick) of Beacon Formation sandstones forming sandstone type interlayers within the Ferrar Dolerite sills.
  - BEACON SUPERGROUP**  
Mainly fine-grained, cross-bedded, coarse to medium-grained sandstone with a lepidolite to quartzose composition. Minor intercalations of conglomerate, black shale, carbonaceous or noncarbonaceous siltstone and minor coal occur as well. A Permian (P) to Upper Triassic age is inferable from "Glossopertis flora" at Beta Peak and Dronning Maud Bay.

- TERRANES OF THE ROSS OROGEN**
- WILSON TERRANE**
  - GRANITE HARBOUR IGNEOUS COMPLEX**
  - WILSON METAMORPHIC COMPLEX**

- GRANITE HARBOUR GRANODIORITE AND GRANITE (GhGr)**  
Granite and granodiorite intruded in the Wilson metamorphic complex below 4000m (see contour lines). At some places, an intrusive breccia forms large and thick bodies made up of migmatite gneiss blocks and pods in a granitoid matrix.
- AMPHIBOLITE FACIES METASEDIMENTS (Ww)**  
From to meso-epizone facies and to intercalations of quartz-biotite metasediments. Biotite cooling ages indicate that metamorphism is older than 400 Ma. (Only in the geological cross-section A-A').

- SYMBOLS**
- Geological boundary.
  - Bedding.
  - Magnetic fluidity.
  - Felsic and mafic dyke swarms. A lot of dykes, both felsic and mafic in composition, belong to the Granite Harbour Igneous Complex, but several mafic dykes pertain to the McMurdo Igneous Complex.
  - Fossils (branchiopods).
  - Fossil plants.
  - Fault: tick (on the downthrown side) or arrows where the sense of motion is known.
  - Shots ACRUP 1993-1994.



**PREVIOUS WORK**

The region considered here was first visited during the Heroic Age of Antarctic exploration. The Magnetic Pole Party of the Shackleton Expedition (1907-1909) crossed the Transantarctic Mountains at the David Glacier. The party included two geologists, Edgeworth David and Douglas Mawson, who did geological observations during the sleeping toward the Magnetic Pole. The party also collected fossils from the Beacon Supergroup. The party also collected fossils from the Beacon Supergroup. The party also collected fossils from the Beacon Supergroup.

**SHORT DESCRIPTION OF GEOLOGY**

The Mount Joyce quadrangle encompasses an early Palaeozoic granitic basement and a fast-lying cover spanning in age from Permian to Jurassic. The basement consists of large bodies of the late Cambrian Granite Harbour Igneous Complex, belonging to the Wilson Terrane, the westernmost of the tectonic units which form the Ross Orogen at the Pacific edge of the Transantarctic Mountains. During the Palaeozoic this orogen was uplifted and eroded. On the resulting peneplain surface, the Permian-Triassic Beacon Sandstone was deposited, which in turn was covered by large flows of the Jurassic Kirkpatrick Basalt. The Jurassic Ferrar Dolerite formed sills chiefly along the basal Beacon horizon.

**LITHOSTRATIGRAPHY**

**Wilson Terrane**  
**Granite Harbour Igneous Complex**  
During the last fifteen years, some petrographic and geochemical differences with a regional significance have been described for these plutonic rocks in Victoria Land, particularly in the area between Aviator Glacier and the Mountaineer Range. Mount Murchison quadrangle (Capponi et al., 1997; Borg et al., 1994; Borg et al., 1998) and Vetter and Tessensohn (1987) stated that these intrusive rocks occur in two NW-SE trending belts throughout the Wilson Terrane, which are interpreted as the magmatic signature of an active continental margin. The western belt is made up of S-type, peraluminous, two-mica and many K-feldspar porphyritic granite; conical-type hornblende granodiorite and tonalite form minor plutons and dykes. The eastern belt comprise H-type, mainly granodioritic to tonalitic intrusive rocks. All the granitic bodies that occur in this quadrangle belong to the western belt.

**Granite Harbour Granodiorite and Granite (GhGr)**  
These rocks form the massifs that crop out in easternmost part of the quadrangle, i.e. in the Prince Albert Mountains north and south of the David Glacier (Mount Priestley, Mount Stephen, Mount Nunatak, and Mount George Murray). Intrusive rocks mainly consist of coarse-grained disacrystic granites and granodiorites, with K-feldspar megacrysts. Microgranular mafic enclaves of dioritic composition are widely diffuse in the granodiorite, while their abundance strongly decreases in the granite. The latter is characterized by a foliated texture defined by elongated K-feldspar megacrysts and mafic enclaves; the latter have strongly elongated shapes parallel to the K-feldspar alignment. This fabric corresponds to a magmatic foliation, well developed at Mount George Murray. The granites and granodiorites are characterized by a foliated texture defined by elongated K-feldspar megacrysts, show poorly foliated or unfoliated fabrics and are further characterised by diffuse occurrence of megacrystic dykes of variable thickness.

**Beacon Supergroup**  
The base of the Beacon Supergroup is represented by a remarkable peneplain surface, which is equivalent to the Kuku Peneplain as defined in the Dry Valley (Barnett et al., 1980). Above this surface the classic Beacon deposits unconformably rest on the underlying basement. In the Fryberg Mountains quadrangle the Beacon strata rest on the Late Palaeozoic diamictite (50-70 m thick; Collins & Kemp, 1983). The same siltites at the base of the Beacon, have been found in the SW corner of the Mount Melbourne quadrangle. The occurrence of these siltites confirms the pre-Permian erosion of the basement. The "Glossopertis flora" newly discovered at Beta Peak (Pertusati et al., 1999) redefines the presence of the Permian interval of the Beacon succession also in this region. The abundant fossil relics were found in a coal-rich sequence of arenitic to shaly strata cropping out at a middle to large SSE of Beta Peak. The sequence is about 40 m thick and is exposed over some hundred metres of length. Moreover, well preserved *Dicroidium odontopteroides* subsp. *odontoides* and *Dicroidium zuberi* relics to be ascribed to *Lower Triassic* (*Ladinian*) and *Early Triassic* (*Scoythin*) (Pertusati et al., 1999). Brambila (1959) obtained data were found at Benson Knob, southern Ricker Hills. It must be underlined that the occurrence of both "Glossopertis flora" and *Dicroidium* in these sediments implies existence of both the Takouana Formation (Dow and Neel, 1974) and the Section Peak Formation (Collinson et al., 1993), but problems of stratigraphic correlation and nomenclature arise as well as we cannot locate at any place the contact between the two lithostratigraphic units. As a consequence, we have indicated all the Beacon Supergroup outcrops as "Beacon sediments" (Bs), and attributed the index T1 (Takouana Formation) and S2 (Section Peak Formation) only to the Bs outcrops at Beta Peak and Benson Knob (southern Ricker Hills), respectively, where the guide fossils, which occur (Pertusati et al., 1999). The contact between the Beacon sediments and the underlying Ross age granitic basement is exposed in the NE side of the Conroy Range Quadrangle, adjacent to the southern boundary of this map; the Ross basement is directly in contact with the Ferrar Dolerite in the NE corner of the Conroy Range Quadrangle (southern slopes of Mt Fearn).

**Ferrari Dolerite (Fd)**  
Major outcrops of Beacon Supergroup are at Ricker Hills, Thomas Rock, Pudding Bath, Richards Nunatak, the unnamed outcrop west of the Milton, and the northernmost part of the Ford Peak. Minor outcrops occur in the upper parts of Mount Howard and Mount Hill. Several 10 to 100 m-wide slices and bodies of Beacon sandstone are incorporated in the Ferrar dolerites at Mount Joyce (northern side), Mount Howard, the Milton, and Mount Bowen (eastern side). Due to the cross-bedding inside these blocks, and their filling caused by the emplacement of the dolerite sills, the attitude of the Beacon sandstones can not be always correctly estimated. Anyway, at large scale the sandstone beds dip very gently toward south and/or southwest, and the dolerite sills share the same attitude.

**Kirkpatrick Basalt (Kb)**  
The Kirkpatrick Basalt (Kb) consists of amygdaloidal lavas with rare tuffaceous and sedimentary interbeds. The lavas are typical tholeiites with two pyroxenes, plagioclase and glassy mesostasis. Vugs and cavities are very frequent and filled by secondary minerals. In most cases zoilites (Vezzalini et al., 1994). The lowermost part of the sequence was distinguished as a separate unit by Elliot et al. (1984), and named the Exposure Hill Formation by the same authors. It consists of volcaniclastic matrix-supported sediments with elements up to some metres in size; vegetal relics are common, and many sandstones and marls occur at some levels.

**Ricker Hills Tillite**  
Tillite sediments crop out at different elevations in the Ricker Hills area. They are covered by Late Pleistocene glacial drift, and lie conformably on the Beacon sandstone and Ferrar Dolerite. Tillite consists of massive matrix-supported diamictite with frequent stratified pebbles and cobbles. Clasts are mainly dolerite, basalt, sandstone, marl, charcoal and siliceous material. The sandy-silt matrix is yellowish at the weathered surface and grey in the interior. The maximum thickness is 20 m. These tillites can tentatively be correlated to the Sirius Formation in the Dry Valley (Micozzi-Piccoferri?).

**McMurdo Igneous Complex**  
No major outcrop occurs in this quadrangle and the presence of McMurdo volcanic rocks is limited to dyke swarms, emplaced in the Granite Harbour Intrusive rocks. Seemingly, no McMurdo dykes was emplaced in the Ferrar Dolerite and Kirkpatrick Basalt, but due to similar field aspect and colour, the possible occurrence of dykes could go unnoticed.

**ROSS TECTONICS**  
In this quadrangle the effects of the Ross tectonic events are not evident, apart the occurrence itself of the Granite Harbour intrusive rocks which were interpreted as the magmatic signature of a Ross-age continental margin convergence. The most common mesozoic structure is the magmatic foliation, which characterizes most of the intrusions belonging to the Larson Granodiorite. The occurrence of isolated intrusive bodies, testifies that the emplacement of Granite Harbour Igneous Complex was coeval with the Late Cambrian - Early Ordovician Ross Orogen deformation. From a geophysical point of view, the magmatic lineament flanking the eastern side of the Glass Anomaly Complex (Bozzo et al., 1997) separates areas featuring contrasting high-frequency anomalies as well as more regional patterns. This NW-SE magnetic lineament crosses the David Glacier in the NE corner of the quadrangle, and continues in the adjoining Relief Inlet quadrangle, roughly parallel to the features of the Ross Sea (Fitzgerald & Bozzo, 1999) the "Central Victoria Land Deformation". This lineament is tentatively interpreted to be the unexposed southern continuation of the Eyles Thrust of Heroic Age, recognized from geological work and magnetic survey much more northwards, along the Pacific Coast (Pillitteri & Klein Schmidt, 1991; Bosum et al., 1989) and Roland (1991) suggested that this lineament marks the boundary between the Ross Orogen and the East Antarctic shield.

**Post-Ross Tectonics**  
The most striking and widespread feature related to the post-Ross tectonics, is represented by the regional unconformity along the pre-Beacon peneplain surface. This surface is the only record that testifies a pre-Permian uplift and erosion of the Ross Orogen. At the regional scale this surface postdates the emplacement of Devonian Antrim Igneous Complex in the Mount Murchison Quadrangle (Capponi et al., 1997) and predates the deposition of Beacon strata with *Glossopertis flora* of Permian age, cropping out in this quadrangle.

The most prominent topographic and tectonic lineament linked to the post-Ross tectonics is the David Glacier. The northern shoulder of this glacier corresponds to a system of NW-SE trending normal faults steeply dipping toward southwest, with a minimum vertical throw of 200 m. To the south of the David Glacier, the Late Palaeozoic-Mesozoic volcano-sedimentary cover weaves dips (2°-4°) toward south and southwest, and is affected by gently folding at the map scale. Two, NW-SE and NE-SW trending fault systems delimit the Palaeozoic basement and the volcano-sedimentary cover with a vertical throw of some hundred metres on single fault. Their occurrence has been recognised on the basis of: (i) difference in elevation of the Beacon strata along east-west sections, (ii) sub-topography (Delisle, 1994a) and (iii) local tilting (up to 15°) of sedimentary layering (e.g. north of Mt. Bowen and Morris Basin). The NW-SE system is more prominent than the NE-SW one; the latter corresponds to a NW dipping fault that lowered the area west of the Hollingsworth Glacier. On the fault surfaces, which can be rarely observed, the slickensides and striae are down dip with a minor component of strike-slip motion. According to Fitzgerald et al. (1999) and Fitzgerald (1992), the David Glacier Lineament separates two segments of the Transantarctic Mountains characterized by different amount of uplift and direction of the rift shoulder escarpment. Mazzanti et al. (1997) interpreted the David Glacier Lineament as a major transfer structure, separating the northern and southern Victoria Land segments. The data on the distribution of Mesozoic and Cenozoic rocks and of the Kuku Peneplain suggest a smaller amount of uplift for the southern segment with respect to the northern Victoria Land (Fitzgerald, 1992). The David Glacier Lineament also changes in direction of the rift shoulder escarpment of the West Antarctic Rift System, which is locally right-lateral. Emplaced with a distal sense of shearing (Stachewski and Thiede, 1992).

The age and activity of the David Glacier Lineament is not well constrained; its offshore prolongation affects the acoustic basement only and not the Cenozoic sequences. However, a genetic link between the E-W trending features and the Jurassic magmatism is suggested by Wilson (1995) and Bozzo et al. (1997); hence, a Mesozoic-Cenozoic reactivation of these tectonic features is probable. In the case the David Glacier Lineament may well have influenced the tectonic patterns of the moderately uplifted Prince Albert Mountains block during diastrophic phases, which occurred along the margin of the East Antarctic Craton in the Mesozoic and Cenozoic. Other field data by Rossetti and Storti (1998) indicate that the Cenozoic tectonic framework is dominated by NW-SE trending right-lateral strike-slip faulting, which is in good agreement with the features of the Ross Sea. The Mount Joyce and the adjoining Relief Inlet quadrangles were the site of the ACRUP-1 (Antarctic Crustal Profile), a geophysical experiment which complements the previous GANEX V system over the Deep Freeze Range crust (O'Connell and Bozzo, 1993). One of the most interesting results from the onshore part of this seismic experiment is the presence of a 7.7 to 7.8 km/s layer just beneath the interpreted Moho. This could be a cushion of proterozoic mantle material similar to the one detected at the base of Deep Freeze Range crust (O'Connell and Bozzo, 1993). This layer may have a regional extent and be related to the thermal perturbation of the mantle beneath the Transantarctic Mountains. Progressive migration of such a layer beneath the Transantarctic Mountains could be the origin of differential thermal uplift of different crustal blocks (Della Vedova et al., 1997).

**ACKNOWLEDGMENTS**  
The authors are grateful to G. Brambila for paleontological studies of Triassic flora, and to E. Bozzo and F. Ferraccioli for their contribution on the tectonics. G. Di Vincenzo, A. Foresti, C. Ghazze, S. Rocchi and D. Visoni are thanked for providing data on the Palaeozoic magmatism.

- CENOZOIC MAGMATISM AT THE ROSS SEA MARGIN**
- McMurdo Igneous Complex: Meander Granite and Syenite (Mg), Maiti, Hallett and Bellingshore Volcanic Suites (B)
  - Kirkpatrick Basalt (Kb): Ferrar Dolerite and Beacon Supergroup (Bs)
  - Admiralty Igneous Complex
  - Tillites
  - Faults
- TERRANES AND UNITS OF THE ROSS OROGEN**
- Wilson Terrane: Wilson Metamorphic Complex (Ww), Granite Harbour Igneous Complex (Gh)
  - Descent Ridge Unit in the Mountaineer Range and post-orogenic correlative, at the eastern margin of the Larimer Range
  - Bowers Terrane
  - Milton Schist
  - Robertson Bay Terrane

- SYNOPTIC TABLE OF STRATIGRAPHIC AND TECTONIC RELATIONSHIPS IN NORTHERN VICTORIA LAND**
- Units occurring in the quadrangle are highlighted in yellow
- McMurdo Igneous Complex: Cenozoic
  - Wilson Terrane: Devonian - Carboniferous
  - Granite Harbour Igneous Complex: Devonian - Carboniferous
  - Wilson Metamorphic Complex: Devonian - Carboniferous
  - Descent Ridge Unit: Devonian - Carboniferous
  - Bowers Terrane: Devonian - Carboniferous
  - Milton Schist: Devonian - Carboniferous
  - Robertson Bay Terrane: Devonian - Carboniferous
  - Beacon Supergroup: Permian - Triassic
  - Ferrar Dolerite: Permian - Triassic
  - Kirkpatrick Basalt: Jurassic
  - McMurdo Igneous Complex: Mesozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic
  - Descent Ridge Unit: Palaeozoic
  - Bowers Terrane: Palaeozoic
  - Milton Schist: Palaeozoic
  - Robertson Bay Terrane: Palaeozoic
  - Beacon Supergroup: Palaeozoic
  - Ferrar Dolerite: Palaeozoic
  - Kirkpatrick Basalt: Palaeozoic
  - McMurdo Igneous Complex: Palaeozoic
  - Wilson Terrane: Palaeozoic
  - Wilson Metamorphic Complex: Palaeozoic